

Particle Acceleration in Magnetically Dominated Turbulence

Luca Comisso


Department of Astronomy, Columbia University
Columbia Astrophysics Laboratory, Columbia University

Princeton Center for Theoretical Sciences
October 30, 2020




Some Prominent Non-Thermal Distributions


Cosmic Ray Spectra of Various Experiments




Pulsar Wind Nebulae (Meyer *et al.* 2010)





Blazars (Abdo *et al.* 2011)



Examples of Turbulent Plasmas



Turbulence Cascade from Fully-Kinetic PIC Simulations




Turbulence develops from uncorrelated initial fluctuations $\delta \mathbf{B}$

Mean magnetic field $\langle \mathbf{B} \rangle = B_0 \hat{\mathbf{z}}$


e^-e^+ plasma with $\sigma_0 = \delta B_{\text{rms}0}^2 / 4\pi h_0 \gg 1$

$$d_{e0} = \left(\frac{\gamma_{th} m_e c^2}{4\pi n e^2} \right)^{1/2}, \quad \frac{L}{d_{e0}} = 820 \Rightarrow 2460^3 \text{ cells}$$

Copious Formation of Current Sheets




Self-Consistent Development of Reconnection Layers





The large computational domain allows the turbulent cascade to develop reconnection layers with flux ropes/plasmoids

Reconnection with flux ropes in dedicated lab experiment



Gekelman *et al.* 2016

Conditions for Triggering Plasmoid-Mediated Recon.



Comisso & Sironi 2019

- Obtain relativistic tearing growth rate

$$\gamma^{1/2} \tau_H^{1/2} \left(\frac{\lambda}{d_e} \right)^{3/2} \frac{\Gamma[(\Upsilon - 1)/4]}{\Gamma[(\Upsilon + 5)/4]} = -\frac{8}{\pi} \Delta'$$

$$\tau_H = \frac{1}{k_\xi v_{A\lambda}}, \quad \Upsilon = \gamma \tau_H \frac{\lambda}{d_e},$$


$$d_e = \left(\frac{\gamma_{th} m_e c^2}{4\pi n e^2} \right)^{1/2}$$

- Use principle of least time (Comisso *et al.* 2016) to obtain the conditions


$$\frac{\lambda}{\xi} \simeq \left(\frac{d_e}{\xi} \right)^{2/3} \left[\ln \left(\frac{1}{\sqrt{\varepsilon}} \left(\frac{d_e}{\xi} \right)^{\frac{4-\alpha}{6}} \right) \right]^{-1/3}$$

$$\gamma \frac{\xi}{v_{A\lambda}} \simeq \ln \left(\frac{1}{\sqrt{\varepsilon}} \left(\frac{d_e}{\xi} \right)^{\frac{4-\alpha}{6}} \right) \gg 1$$

Reconnecting Current Sheets Energize Particles





Development of Non-Thermal Particles




- Turbulence+Reconnection generate a nonthermal particle spectrum with a power-law energy range

Comisso & Sironi 2018, 2019





See also Zhdankin, Nättilä, Wong, *et al.*









How are Turbulence,
Reconnection, and
Particle Acceleration
Interconnected?


Energization History for a Typical Particle



Energization History for Another Typical Particle





Two Stages of Particle Acceleration



- ▶ Particles belonging to the non-thermal tail experience a sudden energy jump from $\gamma \sim \gamma_{th}$ to $\gamma \sim \sigma_0 \gamma_{th} \gg \gamma_{th}$
- ▶ Particle continue to gain energy with a slower rate from $\gamma \sim \sigma_0 \gamma_{th}$ to much higher energies (up to $\gamma \sim \gamma_{cutoff}$).




Particle Injection at Reconnecting Current Sheets



Comisso & Sironi 2018, 2019:

- ▶ At injection, most of the particles reside at $|J_{z,p}| \geq 2 J_{z,\text{rms}}$
- ▶ Locations with $|J_{z,p}| \geq 2 J_{z,\text{rms}}$ are strong current sheets



$$(W_{\parallel,\perp}(t) = q \int_0^t \mathbf{E}_{\parallel,\perp}(t') \cdot \mathbf{v}(t') dt')$$



- ▶ $\Delta\gamma_{\text{inj}} \sim W_{\parallel}/m_e c^2 \sim \sigma_0 \gamma_{th0}$
- ▶ $\mathbf{v} \cdot \mathbf{E}_{\parallel}$ energization is important initially (low $\Delta\gamma$ -range)
- ▶ $\mathbf{v} \cdot \mathbf{E}_{\perp}$ energization gives the majority of particle energy

Comisso & Sironi 2019


Stochastic Particle Acceleration



- ▶ From the self-consistent particles evolution we find (for $\gamma_* \gtrsim \gamma_\sigma$):

$$\langle(\Delta\gamma)^2\rangle \propto \Delta t \Rightarrow \text{Diffusive behavior in energy space}$$
- ▶ The energy diffusion coefficient (in units of c/l) can be calculated as $D_\gamma = \frac{\langle(\Delta\gamma)^2\rangle}{2\Delta t}$ (see also Wong et al 20 for similar PIC results)
- ▶ The power-law tail of the particle spectrum starts at $\gamma/\gamma_\sigma \gtrsim 1$


Two-Stage Acceleration Process



Comisso & Sironi 2018, 2019

- ▶ Injection phase controlled by $E_{\parallel} \Rightarrow \frac{d\langle \gamma \rangle}{dt} = \frac{e}{mc} \beta_R \delta B_{\text{rms}}$
- ▶ Stochastic Acc. controlled by $D_{\gamma} \Rightarrow \frac{d\langle \gamma \rangle}{dt} = 0.4 \sigma \left(\frac{c}{l} \right) \gamma$


The Puzzling Radio Spectrum of the Crab Nebula



Zanin 2017, Lyutikov *et al.* 2019




- Why the spectrum in the radio is so hard ($\nu F_\nu \propto \nu^{0.7}$) ?
- Isotropic distribution of electrons implies $dN/d\gamma \propto \gamma^{-1.6}$
- An anisotropic distribution modifies this “requirement”

Anisotropy of the Pitch Angle Distribution



- ▶ Here: $\cos \alpha = \frac{\mathbf{v} \cdot \mathbf{B}}{|\mathbf{v}| |\mathbf{B}|}$, $\int_{-1}^1 f(\cos \alpha, \gamma) d(\cos \alpha) = 1$
- ▶ The two stage acceleration process leads to anisotropic pitch angle distributions.

Hardened Synchrotron Spectrum




- ▶ Low frequencies:
 $\nu F_\nu \propto \nu^{4/3}$
- ▶ Intermediate frequencies:
 $\nu F_\nu \propto \nu^{(3-p+2q)/(2+q)}$
- ▶ High frequencies:
 $\nu F_\nu \propto \nu^{(3-p)/2}$

Comisso, Sobacchi, Sironi 2020

- ▶ Fully Kinetic Simultaneous Treatment of
“Turbulence + Reconnection”
- ▶ High-Energy Particles are Generated
Self-Consistently as a By-Product of
Turbulence + Reconnection
- ▶ Particle Acceleration Follows a Two-Stage Process:
Acceleration from Reconnection followed by
Stochastic Acceleration by scattering off
Turbulent Fluctuations
- ▶ Turbulence + Reconnection Generate Anisotropic
Pitch Angle Distributions
(Important for the Synchrotron Emission)


The anisotropy ranges are magnetization dependent



Comisso & Sironi 2019

- ▶ $\gamma \sim (\sigma_0/2)\gamma_{th0} \Rightarrow$ particles $\mathbf{v}$ are strongly anti/aligned with $\mathbf{B}$
- ▶ $\gamma \sim 5(\sigma_0/2)\gamma_{th0} \Rightarrow$ distribution with minima at $\cos \alpha = \pm 1, 0$
- ▶ $\gamma \gg 5(\sigma_0/2)\gamma_{th0} \Rightarrow$ particles $\mathbf{v}$ are mostly perpendicular to $\mathbf{B}$

What if we artificially remove $E_{\parallel}$?



- ▶ The normalization drops by 2 orders of magnitude (only $\sim 0.2\%$ of the particles in the non-thermal tail)
- ▶ On the other hand, the slope p is similar, as the high-energy particles “forget” their initial conditions

Biased Random Walk in Momentum Space

Fokker-Planck equation in energy space:

$$\frac{\partial N}{\partial t} = -\frac{\partial}{\partial \gamma} (A_\gamma N) + \frac{\partial^2}{\partial \gamma^2} (D_\gamma N)$$

The convection coefficient A_γ is related to the diffusion coefficient in energy space as

$$A_\gamma = \frac{d\langle \gamma \rangle}{dt} = \frac{1}{\gamma^2} \frac{\partial}{\partial \gamma} (\gamma^2 D_\gamma)$$

Using the previously obtained $D_\gamma = 0.1\sigma \left(\frac{c}{l}\right) \gamma^2$ we get

$$\frac{d\langle \gamma \rangle}{dt} = 0.4\sigma \left(\frac{c}{l}\right) \gamma$$